



Dale and Betty Bumpers

VACCINE RESEARCH CENTER

National Institute of Allergy and Infectious Diseases
National Institutes of Health

Clinical Trials of HIV Vaccines

Barney S. Graham, M.D., Ph.D.
Chief, Clinical Trials Core and Viral Pathogenesis Laboratory
Vaccine Research Center/NIAID/NIH
40 Convent Drive, MSC 3017
Building 40, Room 2502
Bethesda, MD 20892-3017
Phone: (301) 594-8468
FAX: (301) 480-2771
e-mail: bgraham@nih.gov

Introduction

The HIV-1 pandemic has grown to become one of the greatest infectious disease threats to human health and social stability that the world has ever encountered. Nearly 40 million persons are living with HIV-1 infection and more than 21 million people have already died from HIV-induced disease. Although effective anti-retroviral therapy has slowed the epidemic in some industrialized countries, worldwide there are still an estimated 15,000 new HIV infections occurring daily. In addition to the vast personal suffering, the loss of young adult parents, caretakers, and wage-earners, HIV has created an unprecedented strain on the social and economic infrastructure of many developing countries, particularly in Sub-Saharan Africa. These facts make it imperative that the epidemic be controlled as rapidly as possible through prevention of new infections. Although education and available public health approaches should be vigorously pursued, development of a preventive vaccine is the best hope of controlling the HIV epidemic.

New molecular tools in virology and immunology, new adjuvants, new gene expression systems, new antigen delivery systems, recent discoveries in HIV entry and pathogenesis, evidence that natural immunity is achieved in rare instances, and promising studies of candidate vaccines in animal models have provided reasons to hope that developing a safe and effective AIDS vaccine will be possible. However, some have argued that preventive vaccination for AIDS will not be possible (1), and the complex biology of HIV-1 makes this a daunting task.

Expectations of a Vaccine Against HIV

Vaccines developed during the last century have provided unprecedented health and freedom from epidemics of many previously common infectious diseases. They have worked by protecting the vaccinated individual from the consequences of infection, but also by reducing the incidence of transmission within the population, diminishing the spread of epidemics. The ultimate goal for HIV vaccine development is to find an approach that can prevent infection in an exposed individual, or lead to rapid clearance of infected cells to avoid persistent infection. However, no current licensed vaccine for other viral pathogens is known to fully prevent infection, and most are effective because they limit the replication and spread of the pathogen below the threshold for clinical expression of disease. It is unlikely that vaccine-induced immune responses will be able to prevent the establishment of latency. It has been shown that a small proportion of infected CD4⁺ T cells become quiescent, allowing viral latency to be established in a reservoir of long-lived cells (2). Therefore, a more realistic initial goal for HIV vaccine development is to achieve a dampening of the initial viremia in an infected individual, maintenance of a low virus load, and prevention of progression to AIDS. Altering the disease course in individuals could potentially have a large impact on the spread of HIV within a population. The determinants of epidemic spread can be expressed as $R_0 = \beta \times c \times D$, where R_0 is the reproductive rate of the epidemic or a measure of spread; β is the transmission efficiency of the agent; c is the frequency of new partners or new transmission opportunities; and D is the duration of transmissibility. If $R_0 > 1$ the epidemic will spread, and if $R_0 < 1$ the epidemic will

diminish (3). Effective vaccination has the potential to change both β and D , while education, surveillance, and traditional public health approaches can alter “ c ”. The significant impact on the incidence of new infections from the use of highly active anti-retroviral treatment (HAART) and natural history studies encourage the idea that having vaccine-induced immune responses that maintain a low viral load in infected individuals will reduce transmission efficiency in the population (4). Therefore, the initial aim in vaccine development is to identify approaches that will induce immune responses that control infection and prevent disease in individuals, and slow the epidemic spread of HIV within a population.

Assumptions Relevant to HIV Vaccine Design

Vaccine-induced immunity is possible. While there are many challenges remaining, there are several observations that suggest vaccine development is feasible. First, HIV transmission is relatively inefficient. On average more than 200 exposures are required to cause one infection in settings of sexual transmission (5) or needle stick injuries (6). Therefore, modest improvement in anti-viral defenses may have a profound impact on the transmissibility of HIV. Furthermore, based on analysis of HIV isolates in acute infection, most individuals are infected with a very small number of infectious particles, and in many cases a single virion (7,8). The small inoculum size improves the chances that vaccine-induced immunity could prevent infection. In addition, these data suggest that transmitted virus may have limited structural and genotypic features, which would further improve the chances for identifying mechanisms of protective immunity. Secondly, there are examples of natural immunity from studies of highly-exposed, uninfected

(9,10) and long-term nonprogressor (11) populations. There is also evidence from western Africa, where there are concurrent epidemics of HIV-1 and HIV-2, that prior infection with the less virulent HIV-2 confers some protection against HIV-1 infection (12). Finally, there are now many examples of passive protection and vaccine-induced immunity in nonhuman primate models of lentivirus infection.

Timing of the HIV-specific immune response is critical. There are several parameters of the vaccine-induced immune response that will determine its ability to protect the host from infection or disease including specificity, functional properties, magnitude, and compartmentalization. Another critical factor is timing. The timing of the immune response with respect to initial virus infection and spread is particularly important in the case of HIV-1 infection. One reason for this is that the longer HIV-1 replicates in the host, the more diverse variants evolve that may allow escape from subsequent immune responses. In addition, once HIV-1 resides in the extracellular space of lymph node germinal centers and in latently infected cellular reservoirs, or is sequestered in the central nervous system and other sites that are relatively protected from immune responses, it is unlikely that it can be fully eliminated from the host. After the initial burst of virus replication and high-titer viremia the titer of virus in plasma is reduced to a lower level by the initial immune responses and establishes a new plateau about 6 months after infection, referred to as the viral load “set point”. The immune response to HIV-1 infection includes a number of potent effector responses that at best achieve a steady state in which virus clearance matches virus production (13). The magnitude of the viral load “set point” correlates with the rate of immune system destruction (14). The important advantage of vaccine-

induced immune responses is that they are induced prior to infection and can be recalled more rapidly than primary effector mechanisms. Therefore, the success of vaccination may hinge upon altering events that occur in the early hours following HIV-1 exposure.

Vaccines work through induction of adaptive immune responses. Preventive vaccines work through establishing immunologic memory for antigenic structures presented by the pathogen or by infected cells. Therefore, the immunologic “tool box” accessible for vaccine-induced immunity only includes elements of the adaptive immune response. The basic cellular elements of adaptive immunity include the B and T lymphocytes. The primary effector mechanisms important for protection against viruses are antibodies produced by B cells and cytotoxic activity mediated primarily by CD8⁺ T cells. In addition, there are soluble factors produced by activated CD4⁺ and CD8⁺ T cells that have anti-viral activity and can influence the differentiation, expansion, and duration of T cell responses. Elements of the nonadaptive immune system are important during the initial phases of antigen presentation and development of the cytokine microenvironment, mediating many of the activities induced by adjuvants. However, immunity against subsequent infection will be determined by adaptive immune responses with memory for key antigens and functional effector activities that can neutralize the pathogen and rapidly eliminate infected cells.

Neutralizing antibody and cytotoxic T cells are the major effectors of anti-viral immunity.

The correlates of immunity against HIV-1 have not been defined in an absolute sense, but much is known about HIV-specific immune responses associated with long-term survival and

maintenance of low viral loads (13). In addition, there is a general understanding about how different elements of the adaptive immune response should work and these concepts can be tested against observations made in studies of the natural history of HIV infection in humans or experimental data from animal models (15). Alternative vaccine-inducible effector mechanisms mediated by chemokines and other soluble factors produced by T cells may ultimately be shown to have a role in protection (16,17) but in this review I will focus on classical neutralizing antibody and CD8+ cytotoxic T cell activities. There is often debate and speculation about which component of the adaptive immune system is most important for immunity. However, there is abundant evidence for HIV and other virus infections that both antibody and CD8+ CTL are important and perform complementary roles in protection from and control of infection. CD4+ T cells are also of obvious importance, especially for influencing differentiation patterns and expansion of selected lymphocyte populations, but their role as a direct effector of virus clearance is less clear. Therefore, another assumption is that CD4+ T cells will be induced in the process of achieving the appropriate antibody and CD8+ CTL responses, and will not be specifically addressed in this paper.

Antibody is the only component of the adaptive immune response that can neutralize a virus particle prior to infection of a cell and is the only immune response associated with protection for any currently licensed vaccines. Antibody titers can be sustained at high levels in serum and in mucosal secretions and be present at the time of infection. This is unlike T cells that only recognize virus in the context of an already infected cell, and require a few days for activation and expansion of memory populations to respond. Therefore, an effective neutralizing antibody

response will be a critical component of vaccine induced immunity, because it can prevent infection and thereby reduce inoculum size and establishment of latently infected cells.

Neutralization is defined as the ability to reduce infectivity of cell-free virus usually measured in susceptible cells in culture. While this aspect of antibody activity is thought to be the key function associated with protection from infection. There is some debate about the mechanism of neutralization. The identification of specific neutralizing epitopes suggests the site of antibody binding is important. However, it has also been suggested that neutralization occurs when a threshold level of the virion surface is covered by antibody that binds the native envelope oligomer regardless of specificity (18). In either case, it is clear that T cell line-adapted viruses are more susceptible to neutralization than primary field isolates which poses a major hurdle for achieving this immunologic endpoint (19,20).

T cells recognize virus infected cells by specific interactions between the T cell receptor and 8-10 amino acid peptides processed from viral antigens and presented in the context of major histocompatibility complex (MHC) molecules. Therefore, T cells can only recognize and clear virus after infection has occurred. The recognition is restricted by the MHC molecule, which means that the particular epitopes recognized by a given individual will depend on the set of inherited alleles encoding the MHC molecules. While each person should have the capacity to recognize multiple epitopes among the antigens included in HIV-1, the hierarchy of recognition or epitope dominance may vary even among individuals who share MHC haplotypes. These issues suggest that the epitope repertoire in a vaccine will need to have enough breadth to encompass all the relevant MHC haplotypes of potential vaccinees. In addition, it will be

important to induce a broad response in each individual against several viral antigens to diminish the possibility of immune escape through genetic variation and to allow for host selection of dominant epitopes.

The need for CD4⁺ T lymphocytes to initiate the adaptive immune response presents a dilemma since these cells are the major targets for HIV-1 infection. The problem is how to effectively induce protective immunity against HIV-1 without putting vaccine-induced HIV-specific CD4⁺ T cells at risk of infection. This emphasizes the need for effective immune responses, preexistent at the time of HIV exposure, so that virus clearance can be accomplished before the burden of infected cells is sufficient to maintain persistent infection. While CD4⁺ T cells may have some capacity for lysis of HIV-infected cells (20) and production of anti-viral cytokines, the major role is in shaping the immune response by establishing a microenvironment with a particular cytokine composition. For HIV and most other viruses, induction of a Type 1 cytokine profile (production of IL-12, IL-2, and IFN- γ) is more likely to provide protection than induction of Type 2 cytokines (IL-4, IL-5, IL-13). Initial priming with vectors and the use of adjuvants other than alum (which promotes Type 2 responses) would provide an advantage in this regard.

CD8⁺ T cells are the principal effector mechanism of the adaptive immune response to clear virus-infected cells. The CD8⁺ lymphocyte recognizes a virus-infected cell through a cognate interaction between the T cell receptor and a processed peptide epitope presented in the groove of a MHC class I molecule. The lysis of the infected cells occurs through the production and secretion of perforin and granzymes that penetrate the target cell membrane and induce

apoptosis. FasL is also upregulated on the activated CD8⁺ T cell, and can bind Fas on the target cell providing another avenue for inducing apoptosis of the infected target cell. CD8⁺ T cells also produce cytokines with anti-viral properties like IFN- γ and TNF- α , in addition to other soluble factors that may play a role in virus inhibition. The T cell response causes cytopathology not only of the virus-infected cell, but also to a varying degree in bystander cells. This again points to the importance of clearing virus rapidly to diminish the overall cytopathology and illness associated with the immune response to infection.

Data from Studies in Humans and Animal Models

Antibody can prevent HIV infection. It has been directly proven using passive antibody studies in nonhuman primate models of lentivirus infection that sufficient levels of neutralizing antibody can prevent infection. Studies evaluating polyclonal anti-HIV-1 antiserum (22) or monoclonal anti-V3 antibody in HIV-1 infected chimpanzees (23) or polyclonal serum in SIV-infected macaques (24) have shown that when sufficiently high antibody titers are present prior to intravenous challenge that lentivirus infection can be prevented. Importantly, antibody-mediated protection has also been demonstrated against SHIV (a chimeric virus composed of a HIV envelope and SIV nucleocapsid and replication machinery) with an envelope glycoprotein derived from a dual tropic primary HIV isolate, and the protection could be correlated with in vitro neutralizing activity (25). More recently, passive prophylaxis using HIV immune globulin combined with two monoclonal antibodies has protected macaques from vaginal challenge with SHIV (26), and a mixture of three neutralizing monoclonal IgG1 antibodies given to pregnant

macaques has protected their infants from SHIV oral challenge (27). Definitive evidence of antibody-mediated protection in studies of active immunization has been more difficult to demonstrate, but there is an example from early studies performed with whole inactivated SIV vaccines that is provocative. In these studies it was shown that antibodies to cell constituents incorporated into virions during production of challenge stocks were the best correlate of protection. When the virus used to produce vaccine was grown in human cells, and the virus challenge stock was grown in the same human cells, allogenic responses to the human proteins incorporated by the virus were the dominant mechanism of protection (28-31). Studies done with vaccine produced in monkey cells did not show consistent protection. Even though the antibody response was not specific for virus-encoded antigens, this represents an example of vaccine-induced antibody-mediated protection suggesting that protection through induction of virus-specific antibodies may be achievable. When SIV immune globulin was given one day after intravenous challenge with SIV, infection was not prevented, but disease progression was delayed in some animals (32). This again illustrates that the timing of immune responses are critical to the outcome of infection and that preexisting immunity gives the host a distinct advantage.

T cells can control HIV infection. Control of the initial viremia associated with primary HIV infection temporally correlates with the appearance of CD8⁺ cytotoxic T lymphocytes (33,34), and mutations in specific CTL epitopes can be detected in the residual virus population (reviewed in 35). In addition, HIV-specific CD8⁺ CTL activity has been demonstrated in a small subset of uninfected, seronegative commercial sex workers in The Gambia and in Kenya

suggesting transient infection may have occurred inducing protective immunity mediated by CD8⁺ CTL (9,10). In persons who remain uninfected despite significant occupational exposure to HIV-1 contaminated material, studies have also focused on HIV-specific T cell responses. Although HIV-specific antibodies cannot be detected, PBMCs show lymphoproliferative activity when stimulated with HIV-specific peptides (36). HIV-specific CTL responses have also been seen in this cohort (37), suggesting that transient infection may have occurred and been cleared with natural immune defenses. Another subset of persons infected with HIV-1 has persistent infection, but do not progress to AIDS for greater than 12 years. Some of these individuals are infected with virus isolates that replicate poorly (38,39). However, others are infected with viruses that have normal replication capacity, but have maintained a strong and broad set of humoral and cellular HIV-specific immune responses that appears to be responsible for their delayed disease progression. This has been associated with HIV-specific CD4⁺ T cell proliferation (40), strong CD8⁺ CTL activity against multiple epitopes (41), or CD8⁺ CTL responses against selected dominant epitopes (42). Another clue to the importance of T cell responses in the control of HIV has come from the evaluation of HIV-infected persons treated with highly active anti-retroviral therapy (HAART) soon after primary infection. When these persons undergo structured treatment interruptions there is a transient rise in the virus load that results in a boost of functional T cell activity and subsequent control of virus load without HAART (39).

The most compelling evidence for the importance of CD8⁺ CTL for controlling lentivirus infection comes from studies of pathogenesis and vaccine evaluation in nonhuman primate

models. The CD8⁺ CTL response is the best correlate of viremia control after primary SIV infection in macaques (44). There are now several studies using nucleic acid or other recombinant vector approaches that have demonstrated induction of CD8⁺ CTL responses with a weak or absent antibody response does not protect from lentivirus infection, but reduces viral load and delays disease progression. One of the early demonstrations of this was in macaques immunized with recombinant MVA (modified vaccinia Ankara) prior to challenge of macaques with SIV. Vaccination did not prevent infection, and the CTL cell response was associated with delayed disease progression (45). Subsequent studies have shown similar patterns (46-54). As approaches are taken to optimize the CD8⁺ CTL response, such as the addition of an IL-2 adjuvant to a recombinant DNA vaccine regimen (53) or combining modalities of DNA and MVA (54), nearly complete control of subsequent SHIV infection can be achieved. These data are consistent with the premise that vaccines able to establish a preexisting expanded population of HIV-specific CD8⁺ CTL, are likely to delay disease progression in HIV-infected persons.

Clinical Trials of Candidate HIV Vaccines

Overview of concepts evaluated. Clinical trials in seronegative volunteers have been performed to evaluate the safety and immunogenicity of candidate AIDS vaccines in more than 4000 subjects. Several recombinant envelope products, rgp120 or rgp160, produced in insect, yeast or mammalian cells formulated with a variety of adjuvants have been evaluated in clinical trials. Peptides tested to date have been derived from envelope V3 loop or gag sequences of clade B or multiple clades. They have been presented conjugated to an oligolysine backbone, as a

lipopeptide conjugate, mixed with adjuvant, or as a fusion protein with the self-assembling yeast protein Ty as a particle. They have been administered intramuscularly in the deltoid or anterior thigh (to target lymph nodes that also drain the rectal mucosa), rectally and orally as Ty-gag virus-like particles, and orally encapsulated in polylactide co-polymers. Live recombinant vectors including vaccinia, canarypox, and salmonella have been evaluated as well as nucleic acid based vaccines. These vectors have been delivered by a variety of routes and have been constructed to express either single or multiple HIV-1 antigens, from both structural and nonstructural proteins. New trials utilizing recombinant replication-incompetent adenovirus and modified vaccinia Ankara (MVA) are just underway, as well as a novel approach in which tat is the vaccine antigen. Tat is secreted from HIV-infected cells and has a variety of adverse effects on neighboring cells. It is hypothesized that blocking these effects with vaccine-induced antibody will facilitate virus clearance (55). In addition, there have been studies evaluating schedule of administration and combination approaches using more than one product in the immunization regimen (56-57), These studies are listed in the posted Table and referenced when possible. Fortunately, there have been no significant safety concerns other than unacceptable local reactogenicity associated with a few selected adjuvants (58).

Vaccine-induced antibody responses in clinical trials. Neutralizing antibody responses have been induced by immunization with recombinant envelope glycoproteins alone or in combination with poxvirus vectors. The antibody response to immunization with rgp120 alone is in general maximal after the third or fourth injection, is dose-dependent, and can be attenuated unless there is a several month interval between injections. Serum antibody titers have a relatively short half-

life, and while they can be boosted, the titers generally achieve their peak level after the third or fourth injections. Repeated boosting does not prolong the half-life significantly. Therefore, it is likely that recombinant envelope glycoprotein products may find their greatest utility in boosting antibody responses in subjects primed with recombinant vector vaccines (59,60), or other strategies that can induce MHC class I-restricted CTL responses. This combination approach not only adds the CD8+ CTL component to the immune response, but also results in a more durable antibody response. The initial recombinant envelope glycoprotein products were derived from sequences of syncytium-inducing, T cell line-adapted (TCLA), CXCR-4 utilizing X4 viruses from clade B. Newer products, such as the VaxGen B/B product incorporate sequences from primary isolates which utilize CCR5 (R5) combining the rgp120 from HIV-1_{MN} and the rgp120 from HIV-1_{GNR8} (61). Phase I and II studies have defined how parameters of dose, schedule, and formulation affect immunogenicity of purified protein subunit preparations as primary immunogens and as booster immunogens given in combination with other vaccine modalities. The principal findings related to vaccine-induced antibody responses in clinical trials of candidate HIV vaccines are:

- 1) While type-specific neutralization can be induced, particularly to the vaccine antigen, neutralization of typical primary R5 HIV isolates is not induced (62). There are some reports of neutralization of selected R5 HIV strains, but these are strains that are easier to neutralize in general, and how this will translate into protection against more typical neutralization resistant strains is not known.

- 2) Antibody is induced that can bind R5 virus-infected cells (63). It is possible that this represents binding to monomeric envelope determinants, but suggests that the monomeric envelope products currently being tested can produce antibody that recognizes oligomeric envelope structures, even though the affinity and specificity is not sufficient to result in virus neutralization.
- 3) Antigens produced in mammalian cells induce higher titer of neutralizing antibody against TCLA virus than those produced in baculovirus or yeast systems (64-67).
- 4) Recombinant gp120 products induce less binding antibody, but more neutralizing antibody than rgp160 products (66-68). As noted above, the neutralizing activity does not include primary isolate R5 viruses.
- 5) A four dose immunization regimen using envelope glycoprotein is more effective for antibody induction when there is a several month interval between doses (66,67). Intervals of at least 3 to 4 months between the second, third, and fourth immunization increase the magnitude of response.
- 6) A rapid (every month) vaccination schedule using envelope glycoproteins alone results in attenuation of antibody responses after the fourth dose (66). Titers of both binding and neutralizing antibody activities are reduced after a monthly immunization schedule using rgp120 in MF59 (66). The attenuating effect of rapid dosing is not as apparent with other adjuvants (69).
- 7) A fifth dose of rgp120, regardless of interval, does not boost antibody response, but only returns it to previous level.
- 8) The half-life of vaccine antigen-specific antibody titers is <3 months in subjects receiving only rgp120 envelope glycoprotein, regardless of number of doses. The half-life is extended with

gp160 antigens, and is also more prolonged when rgp120 immunization is preceded by priming with poxvirus vectors. The factors underlying antibody maintenance have not been defined.

9) Priming with one subtype and boosting with another demonstrates subtype-specificity in antibody response (70). When a subject is initially immunized with rgp120 derived from a clade B, TCLA X4 HIV strain, subsequent boosting with another clade B strain does not broaden the response significantly, and does not boost the response to the new envelope antigen as well as to the original rgp120.

11) Antigen dose effects on magnitude of antibody production are dependent on the adjuvant formulation. QS21 appears to allow a reduction in the antigen dose by more than 10-100 fold without affecting the magnitude of antibody response (71).

12) Peptide vaccines in general have been weakly immunogenic with one exception. This was a complex peptide that contains T helper epitopes from the C4 domain of gp120 and neutralizing antibody and CTL epitopes from the V3 domain. Peptides from four strains of HIV-1 were combined in incomplete Freund's adjuvant and administered intramuscularly. Neutralizing activity against HIV-1_{MN} was detected in 75% of subjects after the second dose. However, the study was terminated prematurely because of the development of sterile abscesses in a few vaccinees.

13) Recombinant gp160 vaccinia immunization induces an antibody response to the HIV envelope that is slow in developing, often not being detectable until >100 days after inoculation (72),

14) Recombinant vaccinia effectively primes for antibody responses elicited by subsequent boosting with purified recombinant envelope glycoprotein formulations (59,73),

15) Specificity of the antibody response in subject primed with recombinant vaccinia then boosted with recombinant envelop glycoprotein is determined more by the initial antigen expressed by the recombinant vector than by the subsequent envelope antigen given as booster (70,74),

16) Priming immunization with recombinant vaccinia induces type-specific memory for the recombinant gene product that is not boosted by subsequent immunization with heterologous recombinant envelope glycoproteins (70).

17) In subjects immunized with recombinant HIV-1_{LAI} gp160 vaccinia and boosted with HIV-1_{LAI} gp160 produced in baculovirus and formulated with alum, the dominant antibody response in vaccinees was directed against a gp41 epitope (aa 720-740) that was not a major target for antibodies produced by HIV-1_{LAI}-infected persons (75).

18) The HIV-specific antibody response after recombinant canarypox immunization alone is weak, but subsequent boosting with purified recombinant envelope subunit protein induces HIV-specific antibody titers of the same magnitude and quality as 3 or 4 inoculations of the purified recombinant envelope subunit protein alone (76,77).

In summary, neutralizing antibody responses against TCLA viruses induced by the most immunogenic formulations are still 5-10 fold lower than those produced by HIV-1 infection. The responses are type-specific with a relatively short half-life, and are unable to neutralize typical primary isolate R5 viruses.

Vaccine-induced CD8⁺ CTL responses in clinical trials. Induction of HIV-specific CD8⁺ CTL responses generally requires the delivery of vaccine antigens into the cytoplasmic compartment of an antigen presenting cell (APC) for display in a MHC class I molecule on the cell surface. Therefore, vector-based approaches or nucleic acid vaccines that rely on antigen production within the target cell are most effective. Delivering vaccine antigens as purified proteins or even whole inactivated virus will primarily access the endocytic pathway for antigen presentation and lead to CD4⁺ T cell activation. While this is critical for antibody production and important for supporting CD8⁺ CTL development, it is not sufficient for inducing CD8⁺ CTL. In some cases a novel adjuvant or delivery system is able to provide access for these types of vaccines into the cytoplasmic compartment, but in general vector-based vaccines, including nucleic acids, are more potent methods for inducing CD8⁺ CTL. One exception is the use of peptides that incorporate a T cell epitope that can bind directly to an MHC class I molecule on the cell surface and induce CD8⁺ CTL responses. Vector-based vaccines, beginning with recombinant vaccinia products, were first evaluated in clinical trials in the late 1980's with the expressed purpose of achieving vaccine-induced CD8⁺ CTL responses. The induction of CD8⁺ CTL responses has been a primary focus of clinical trials since the mid 1990's.

- 1) It has been found that recombinant vaccinia expressing envelope glycoprotein only, or multiple antigens can consistently induce long-lived CD8⁺ CTL responses in vaccinia-naive subjects (60,78-80).
- 2) HIV-specific CD8⁺ CTL can also be detected in a majority of subjects receiving recombinant canarypoxvirus vectors, and in a subset CTL activity is detectable for >18

months. The activity is at the threshold of detection in classical ^{51}Cr release assays requiring in vitro stimulation and is only detected in 15-30% of subjects at any given time point (81-85). However, unlike antibody responses, vaccine-induced CTL responses are broadly cross-reactive (86). CTLs induced by recombinant canarypox vectors have been shown to lyse target cells infected with primary R5 HIV-1 isolates from multiple clades (86).

- 3) CD8⁺ CTL effectors have also been isolated from rectal mucosa from vaccinees suggesting that T cells induced by parenteral vaccination may provide some level of protection at mucosal surfaces (McElrath et al., unpublished observations).
- 4) Not only is classical MHC class I-restricted cytotoxic activity induced, but vaccine-induced noncytotoxic CD8⁺-mediated suppression of HIV-1 replication has also been demonstrated in recipients of recombinant canarypox vaccines (87).
- 5) Envelope subunits can induce CD4⁺ CTL (21), but rarely induce CD8⁺ CTL even when formulated with novel adjuvants (66). This is an expected result of obligate processing through the endocytic pathway leading to MHC class II presentation. New adjuvants and delivery systems may improve MHC class I presentation of purified protein antigens, but it is unlikely to ever approach the efficiency of antigens produced intracellularly by approaches such as nucleic acid immunization, live recombinant vectors, or live attenuated vaccines.
- 6) In one study with the Ty-gag-VLP without alum CD8⁺ CTL responses were detected in a few subjects (88).

- 7) Subjects injected with a mixture of lipopeptide epitopes formulated with QS21 can induce CD8⁺ CTL responses restricted to selected HLA molecules (89).

In summary, vaccine approaches that are currently being evaluated in clinical trials can induce HIV-specific CD8⁺ CTL activity that is durable and can lyse cells infected with typical primary R5 HIV-1 isolates from multiple clades.

Recent advances in methods to quantitate T cells and evaluate their function are changing the process of vaccine evaluation. Enumeration of functional T cells by IFN- γ ELIspot or intracellular IFN- γ by FACS analysis combined with identification of epitope-specific T cells with MHC-peptide tetramers by FACS analysis (90) have improved the ability to detect vaccine-induced responses by improving sensitivity and reproducibility, allowing the use of cryopreserved cells, and reducing the number of effector cells needed.

Future scientific challenges for HIV vaccine development

Timeline for HIV vaccine development. Although the need for an effective vaccine against HIV is urgent, what is a realistic timeline for identifying the approaches and products needed to achieve vaccine-induced immunity? Some groups initiated a 10-year countdown in 1997 challenging the scientific community to develop an effective HIV vaccine by the year 2007. It is now 2001 and to meet this timeline, the following primary goals will have to be achieved: 1) identification of antigenic structures that can induce neutralizing antibody activity against

primary R5 HIV-1 isolates, 2) development of new approaches to optimize the breadth and magnitude of vaccine-induced memory CTL, and 3) performance of a large scale trial to test the concept that a vaccine-induced CD8⁺ CTL response will modify infection rates or alter the course of naturally acquired HIV infection.

Induction of R5 neutralizing antibody activity. Despite the current optimism there are still many scientific obstacles to overcome in the development of a vaccine for HIV. Most important is the inability to induce broadly cross-reactive neutralizing antibody against typical primary HIV-1 isolates. Many R5 viruses are difficult to neutralize even with high titer, type-specific serum. The relative neutralization resistance of HIV is one of several immune evasion strategies employed by HIV under active investigation. Without a high level of neutralizing activity present at the time of infection, it is unlikely that a vaccine-induced immune response can prevent the establishment of latency and infection of immunoprivileged sites. Empiric approaches are being pursued to find envelope glycoprotein structures that can induce broad neutralizing antibody and include: 1) triggering the envelope glycoprotein into an intermediate structure present during the membrane fusion process, 2) producing envelope glycoproteins with selected mutations that alter glycosylation or folding patterns, 3) production of oligomeric envelope structures, 4) expression of envelope glycoproteins in vectors and virus-like particles, and 5) multi-epitope combinations. In addition, a more systematic approach combining information from structural biology, molecular biology, and epitope mapping to produce novel envelope structures is being pursued. Solving the problem of how to produce an antigenic structure that can induce a neutralizing antibody response effective against typical R5 HIV-1 isolates represents an important step toward

addressing additional questions about the antibody response including: 1) will local induction of mucosal antibody be necessary for protection from HIV infection or will high titer serum antibody be sufficient, 2) can broadly neutralizing antibody be induced with a single antigenic structure, or will mixtures of envelope glycoprotein structures be required, 3) can durable antibody responses be induced, or will repeated booster immunizations be required?

Duration, magnitude, kinetics, and breadth of CTL response. There is evidence from studies in both humans and animal models that a robust CD8⁺ CTL response can control HIV infection. However, the character of those responses and relative degree of protection for a given level and breadth of CTL response have not been defined. There are candidate vaccines currently in Phase II clinical trials that can induce CD8⁺ CTL responses in a portion of subjects. Whether this level of CTL is sufficient for control of HIV infection is not known. However, based on studies in nonhuman primates, it is likely that vaccine induction of a robust CD8⁺ CTL response will lower the viral load set point. The large number of vector-based vaccine delivery approaches, emergence of cytokine adjuvants to specifically direct selected immune responses, and advances in methods used to enumerate T cell responses promise that strategies for consistent vaccine-induction of high magnitude CTL responses are within reach in a time frame of about 5 years. Future studies may need to address the following issues: 1) how broad does the CTL response need to be to control viremia, in terms of number and dominance hierarchy of epitopes, 2) does the CTL response measured in peripheral blood accurately predict control of viremia, or will specific induction of CTL responses in mucosal tissue or lymph nodes be required, 3) can the kinetics of the CTL response or efficiency of killing be influenced by vaccine formulation and

delivery to improve protection, or is the outcome only determined by the magnitude of the response, and 4) will maintenance of sufficient CTL memory require booster immunizations?

The importance of efficacy trial evaluation. One of the next major steps in vaccine development will be the performance of larger scale trials in higher risk populations. Many of the questions involving the importance of HIV genetic variation, mucosal immunity, and duration of vaccine-induced immune responses will be difficult to address until large-scale efficacy trials are implemented. The appropriate timing, vaccine approach, trial design, and trial location for such a study are issues of current controversy and debate. The performance of a Phase III clinical trial should be based on: 1) its potential for defining a biological impact of the vaccine on HIV-induced disease based on results from animal model studies and Phase I/II trials, 2) its potential for answering questions about correlates of immunity, and 3) the importance of establishing a benchmark against which future vaccine design and development can be measured. For example, if a Phase III study can be designed and executed that will show whether CD8⁺ CTL can control HIV viremia, the current vaccine strategies with potential CTL-inducing capacity would be accelerated with a focus on optimizing the magnitude of CD8⁺ induction. If induction of CD8⁺ CTL is not associated with any level of protection, then issues of breadth and compartmentalization of responses will need to be addressed more rigorously in animal models.

In summary, the ultimate vaccine that can prevent persistent HIV-1 infection will probably require a conceptual breakthrough in the understanding of how to elicit broadly neutralizing antibody against primary R5 HIV-1 isolates, and will also involve a number of iterative steps to

achieve optimal HIV-specific CD8+ CTL responses. However, a vaccine aimed at control of viremia, delayed disease progression, and reduced transmission, based on induction of HIV-specific CD8+ CTL could have a significant impact on the AIDS epidemic, and may be within our grasp using currently available technology.

To examine the VRC strategic plan, go to: <http://www.vrc.nih.gov>

Literature Cited

1. Sabin AB. Improbability of effective vaccination against human immunodeficiency virus because of its intracellular transmission and rectal portal of entry. *Proc Natl Acad Sci USA* 1992; 89:8852-5.
2. Sciliciano RF. *Ann Rev Med* 2001; (in press).
3. Holmes KK. Human ecology and behavior and sexually transmitted bacterial infections. *Proc Natl Acad Sci USA* 1994; 91:2448-55.
4. Quinn TC, Wawer MJ, Sewankambo N, Serwadda D, Li C, et al. Viral load and heterosexual transmission of human immunodeficiency virus type 1. Rakai Project Study Group. *N Engl J Med* 2001; 342:921-9.
5. Gray RH, Wawer MJ, Brookmeyer R, Sewankambo NK, Serwadda D, et al. Probability of HIV-1 transmission per coital act in monogamous, heterosexual, HIV-1-discordant couples in Rakai, Uganda. *Lancet* 2001; 357:1149-53.

6. Geberding JL. Incidence and prevalence of human immunodeficiency virus, hepatitis B virus, hepatitis C virus, and cytomegalovirus among health care personnel at risk for blood exposure: final report from a longitudinal study. *J Infect Dis* 1994; 170:1410-7.
7. Wolinsky SM, Wike CM, Korber BTM, Hutto C, Parks WP, et al. Selective transmission of human immunodeficiency virus type-1 variants from mothers to infants. *Science* 1992; 255:1134-7.
8. Wolfs TF, Zwart G, Bakker M, Goudsmit J. HIV-1 genomic RNA diversification following sexual and parenteral virus transmission. *Virology* 1992; 189:103-10.
9. Fowke KR, Nagelkerke NJ, Kimani J, Simonsen JN, Anzala AO, et al. Resistance to HIV-1 infection among persistently seronegative prostitutes in Nairobi, Kenya. *Lancet* 1996; 348:1347-51.
10. Rowland-Jones S, Sutton J, Ariyoshi K, Dong T, Gotch F, et al. HIV-specific cytotoxic T-cells in HIV-exposed but uninfected Gambian women. *Nat Med* 1995; 1:59-64.
11. Wagner R, Leschonsky B, Harrer E, Paulus C, Weber C, et al. Molecular and functional analysis of a conserved CTL epitope in HIV-1 p24 recognized from a long-term nonprogressor: constraints on immune escape associated with targeting a sequence essential for viral replication. *J Immunol* 1999; 162:3727-34.
12. Kokkotou EG, Sankale JL, Mani I, et al. In vitro correlates of HIV-2 mediated HIV-1 protection. *Proc Natl Acad Sci USA* 2000; 97:6797-802.
13. Walker BD. *Ann Rev Med* 2001; (in press).
14. Mellors JW, Rinaldo CR Jr, Gupta P, White RM, Todd JA, et al. Prognosis in HIV-1 infection predicted by the quantity of virus in plasma. *Science* 1996; 272:1167-70.

15. Ada GL. Modern vaccines: the immunological principles of vaccination. *Lancet* 1990; 335:523-6.
16. Walker CM, Moody DJ, Stites DP, Levy JA. CD8+ lymphocytes can control HIV infection in vitro by suppressing virus replication. *Science* 1986; 234:1563-6.
17. Cocchi F, DeVico AL, Garzino-Demo A, Arya SK, Gallo RC, et al. Identification of RANTES, MIP-1 alpha, and MIP-1 beta as the major HIV-suppressive factors produced by CD8+ T cells. *Science* 1995; 270:1811-5.
18. Parren PW, Burton DR. The antiviral activity of antibodies in vitro and in vivo. *Adv Immunol* 2001; 77:195-262.
19. Trkola A, Ketas T, Kewalramani VN, Endorf F, Binley JM, et al. Neutralization sensitivity of human immunodeficiency virus type 1 primary isolates to antibodies and CD4-based reagents is independent of coreceptor usage. *J Virol* 1998; 72:1876-85.
20. Cecilia D, KewalRamani VN, O'Leary J, Volsky B, Nyambi P, et al. Neutralization profiles of primary human immunodeficiency virus type 1 isolates in the context of coreceptor usage. *J Virol* 1998; 72:6988-96.
21. Stanhope PE, Clements ML, Siliciano RF. Human CD4+ cytolytic T lymphocyte responses to a human immunodeficiency virus type 1 gp160 subunit vaccine. *J Infect Dis* 1993; 168:92-100.
22. Prince AM, Reesink H, Pascual D, Horowitz B, Hewlett I, et al. Prevention of HIV infection by passive immunization with HIV immunoglobulin. *AIDS Res Hum Retroviruses* 1991; 7:971-3.

23. Emini EA, Schleif WA, Nunberg JH, Conley AJ, Eda Y, et al. Prevention of HIV-1 infection in chimpanzees by gp120 V3 domain-specific monoclonal antibody. *Nature* 1992; 355:728-30.
24. Putkonen P, Thorstensson R, Ghavamzadeh L, Albert J, Hild K, et al. Prevention of HIV-2 and SIV_{sm} infection by passive immunization in cynomolgus monkeys. *Nature* 1991; 352:436-8.
25. Shibata R, Igarashi T, Haigwood N, Buckler-White A, Ogert R, et al. Neutralizing antibody directed against the HIV-1 envelope glycoprotein can completely block HIV-1/SIV chimeric virus infections of macaque monkeys. *Nat Med* 1999; 5:204-10
26. Mascola JR, Lewis MG, Stiegler G, Harris D, VanCott TC, et al. Protection of macaques against pathogenic simian/human immunodeficiency virus 89.6PD by passive transfer of neutralizing antibodies. *J Virol* 1999; 73:4009-18.
27. Baba TW, Liska V, Hofmann-Lehmann R, Vlasak J, Xu W, et al. Human neutralizing monoclonal antibodies of the IgG1 subtype protect against mucosal simian-human immunodeficiency virus infection. *Nat Med* 2000; 6:200-6.
28. Stott EJ. Anti-cell antibody in macaques. *Nature* 1991; 353:393.
29. Stott EJ, Chan WL, Mills KH, Page M, Taffs F, et al. Preliminary report: protection of cynomolgus macaques against simian immunodeficiency virus by fixed infected-cell vaccine. *Lancet* 1990; 336:1538-41.
30. Langlois AJ, Weinhold KJ, Matthews TJ, Greenberg ML, Bolognesi DP. Detection of anti-human cell antibodies in sera from macaques immunized with whole inactivated virus. *AIDS Res Hum Retrovir* 1992; 8:1641-52

31. Montefiori DC, Cornell RJ, Zhou JY, Zhou JT, Hirsch VM, et al. Complement control proteins, CD46, CD55, and CD59, as common surface constituents of human and simian immunodeficiency viruses and possible targets for vaccine protection. *Virology* 1994; 205:82-92.
32. Haigwood NL, Watson A, Sutton WF, McClure J, Lewis A, et al. Passive immune globulin therapy in the SIV/macaque model: early intervention can alter disease profile. *Immunol Lett* 1996; 51:107-14.
33. Koup RA, Safrit JT, Cao YZ, Andrews CA, McLeod G, et al. Temporal association of cellular immune responses with the initial control of viremia in primary human immunodeficiency virus type 1 syndrome. *J Virol* 1994; 68:4650-55.
34. Borrow P, Lewicki H, Hahn BH, Shaw GM, Oldstone MB. Virus-specific CD8⁺ cytotoxic T-lymphocyte activity associated with control of viremia in primary human immunodeficiency virus type 1 infection. *J Virol* 1994; 68:6103-10.
35. McMichael AJ, Phillips RE. Escape of human immunodeficiency virus from immune control. *Ann Rev Immunol* 1997; 15:271-96.
36. Clerici M, Levin JM, Kessler HA, Harris A, Berzofsky JA, et al. HIV-specific T-helper activity in seronegative health care workers exposed to contaminated blood. *J Am Med Assoc* 1994; 271:42-6.
37. Pinto LA, Sullivan J, Berzofsky JA, Clerici M, Kessler HA, et al. ENV-specific cytotoxic T lymphocyte responses in HIV seronegative health care workers occupationally exposed to HIV-contaminated body fluids. *J Clin Invest* 1995; 96:867-76.

38. Deacon NJ, Tsykin A, Solomon A, Smith K, Ludford-Menting M, et al. Genomic structure of an attenuated quasi species of HIV-1 from blood transfusion donor recipients. *Science* 1995; 270:988-91.
39. Kirchhoff F, Greenough TC, Brettler DB, Sullivan JL, Desrosiers RC. Brief report: Absence of intact nef sequences in a long-term survivor with nonprogressive HIV-1 infection. *N Engl J Med* 1995; 332:228-32.
40. Rosenberg ES, Billingsley JM, Caliendo AM, Boswell SL, Sax PE, et al. Vigorous HIV-1-specific CD4⁺ T cell responses associated with control of viremia. *Science* 1997; 278:1447-5.
41. Ortiz GM, Nixon DF, Trkola A, Binley J, Jin X, et al. HIV-1-specific immune responses in subjects who temporarily contain virus replication after discontinuation of highly active antiretroviral therapy. *J Clin Invest* 1999; 104:R13-8.
42. Goulder PJR, Brander C, Yang Y, Tremblay C, Colbert RA, Addo MM, Rosenberg ES, Nguyen T, Allen R, Trocha A, Altfeld M, He S, Bunce M, Funkhouser R, Pelton SI, Burchett SK, McIntosh K, Korber BTM, Walker BD. Evolution and transmission of stable CTL escape mutations in HIV infection. (in press).
43. Robenberg ES, Altfeld M, Poon SH, Phillips MN, Wilkes BM, et al. Immune control of HIV-1 after early treatment of acute infection. *Nature* 2000; 407:523-6.
44. Reimann KA, Tenner Racz K, Racz P, Montefiori DC, Yasutomi Y, et al. Immunopathogenic events in acute infection of rhesus monkeys with simian immunodeficiency virus of macaques. *J Virol* 1994; 68:2362-70.

45. Hirsch VM, Fuerst TR, Sutter G, Carroll MW, Yang LC, et al. Patterns of viral replication correlate with outcome in simian immunodeficiency virus (SIV)-infected macaques: effect of prior immunization with a trivalent SIV vaccine in modified vaccinia virus Ankara. *J. Virol.* 1996; 70:3741-52.
46. Ourmanov I, Brown CR, Moss B, Carroll M, Wyatt L, et al. Comparative efficacy of recombinant modified vaccinia virus Ankara express simian immunodeficiency virus (SIV) Gag-Pol and/or Env in macaques challenged with pathogenic SIV. *J Virol* 2000; 74:2740-51.
47. Leno M, Kowalski M, Robert-Guroff M. CD8 T cell anti-HIV activity as a complementary protective mechanism in vaccinated chimpanzees. *AIDS* 2000; 14:893-4.
48. Allen TM, Vogel TU, Fuller DH, Mothe BR, Steffen S, et al. Introduction of AIDS virus-specific CTL activity in fresh, unstimulated peripheral blood lymphocytes from rhesus macaques vaccinated with a DNA prime/modified vaccinia virus Ankara boost regimen. *J Immunol* 2000; 164:4968-78.
49. Buge SL, Richardson E, Alipanah S, Markham P, Cheng S, et al. An adenovirus-simian immunodeficiency virus env vaccine elicits humoral, cellular, and mucosal immune responses in rhesus macaques and decreases viral burden following vaginal challenge. *J Virol* 1997; 71:8531-41.
50. Abimiku AG, Robert-Guroff M, Benson J, Tartaglia J, Paoletti E, et al. Long-term survival of SIVmac251-infected macaques previously immunized with NYVAC-SIV vaccines. *J Acq Immun Defic Syndr* 1997; 15:S78-85.

51. Davis NL, Caley IJ, Brown KW, Betts MR, Irlbeck DM, et al. Vaccination of macaques against pathogenic simian immunodeficiency virus with Venezuelan equine encephalitis virus replicon particles. *J Virol* 2000; 74:371-8.
52. Egan MA, Charini WA, Kuroda MJ, Schmitz JE, Racz P, et al. Simian immunodeficiency virus (SIV) gag DNA-vaccinated rhesus monkeys develop secondary cytotoxic T-lymphocyte responses and control viral replication after pathogenic SIV infection. *J Virol* 2000; 74:7485-95.
53. Barouch DH, Santra S, Schmitz JE, Kuroda MJ, Tong-Ming F, et al. Control of viremia and prevention of clinical AIDS in rhesus monkeys by cytokine-augmented DNA vaccination. *Science* 2001; 290:486-92.
54. Amara RR, Villinger F, Altman JD, Lydy SL, O'Neil SP, et al. Control of a mucosal challenge and prevention of AIDS by a multiprotein DNA/MVA vaccine. *Science* 2001; 292:69-74.
55. Cafaro A, Caputo A, Fracasso C, Maggiorella MT, Goletti D, et al. Control of SHIV-89.6P-infection of cynomolgus monkeys by HIV-1 Tat protein vaccine. *Nat Med* 1999; 5:643-50.
56. Graham BS, Karzon DT. AIDS vaccine development. In *Textbook of AIDS Medicine*, ed. TC Merigan, JG Bartlett, D Bolognesi, 1998; 42:689-724. Baltimore: Williams & Wilkins. 2nd ed.
57. Graham BS. Clinical trials of HIV vaccines. In *HIV Molecular Immunology Database Compendium*, ed. BTM Korber, JP Moore, C Brander, BD Walker, BF Haynes, R Koup,

Los Alamos: National Laboratory of Theoretical Biology and Biophysics. 2000; I-20-I-38.

<http://hiv-web.lanl.gov>.

58. Keefer MC, Wolff M, Gorse GG, Graham BS, McElrath MJ, et al. Safety profile of phase I and II preventive HIV-1 vaccination: Experience of the AIDS Vaccine Evaluation Group. *AIDS Res Hum Retrovir* 1997; 13:1163-77.
59. Graham BS, Matthews TJ, Belshe RB, Clements ML, Dolin R, et al. Augmentation of human immunodeficiency virus type 1 neutralizing antibody by priming with gp160 recombinant vaccinia and boosting with rgp160 in vaccinia-naive adults. *J Infect Dis* 1993; 167:533-7.
60. Cooney EL, McElrath MJ, Corey L, Hu SL, Collier AC, et al. Enhanced immunity to human immunodeficiency virus (HIV) envelope elicited by a combined vaccine regimen consisting of priming with a vaccinia recombinant expressing HIV envelope and boosting with gp160 protein. *Proc Natl Acad Sci USA* 1993; 90:1882-6.
61. Berman PW, Huang W, Riddle L, Gray AM, Wrinn T, et al. Development of bivalent (B/E) vaccines able to neutralize CCR5-dependent viruses from the United States and Thailand. *Virology* 1999; 265:1-9.
62. Mascola JR, Snyder SW, Weislow OS, et al. Immunization with envelope subunit vaccine products elicits neutralizing antibodies against laboratory-adapted but not primary isolates of human immunodeficiency virus type 1. *J Infect Dis* 1996; 173:340-8.
63. Gorse GJ, Patel GB, Newman FK, et al. Antibody to native human immunodeficiency virus type 1 envelope glycoproteins induced by IIIB and MN recombinant gp120 vaccines. *Clin Diag Lab Immunol* 1996; 3:378-86.

64. Dolin R, Graham BS, Greenberg SB, et al. The safety and immunogenicity of a human immunodeficiency virus type 1 (HIV-1) recombinant gp160 candidate vaccine in humans. NIAID AIDS Vaccine Clinical Trials Network. *Ann Intern Med* 1991; 114:119-27.
65. Keefer MC, Graham BS, McElrath MJ, et al. Safety and immunogenicity of Env 2-3, a human immunodeficiency virus type 1 candidate vaccine, in combination with a novel adjuvant, MTP- PE/MF59. *AIDS Res Hum Retrovir* 1996; 12:683-93.
66. Graham BS, Keefer MC, McElrath MJ, et al. Safety and immunogenicity of a candidate HIV-1 vaccine in healthy adults: Recombinant glycoprotein (rgp) 120 - A randomized, double-blind trial. *Ann Intern Med* 1996; 125:270-9.
67. Belshe RB, Graham BS, Keefer MC, et al. Neutralizing antibodies to HIV-1 in seronegative volunteers immunized with recombinant gp120 from the MN strain of HIV-1. NIAID AIDS Vaccine Clinical Trials Network. *JAMA* 1994; 272:475-80.
68. Belshe RB, Clements ML, Dolin R, et al. Safety and immunogenicity of a fully glycosylated recombinant gp160 human immunodeficiency virus type 1 vaccine in subjects at low risk of infection. *J Infect Dis* 1993; 168:1387-95.
69. Gorse GJ, Rogers JH, Perry JE, et al. HIV-1 recombinant gp160 vaccine induced antibodies in serum and saliva. *Vaccine* 1995; 13:209-14.
70. Corey L, McElrath MJ, Weinhold K, et al. Cytotoxic T cell and neutralizing antibody responses to HIV-1 envelope with a combination vaccine regimen. *J Infect Dis* 1998; 177:301-9.
71. Evans TG, McElrath MJ, Matthews TJ, Montefiori DC, Wolff M, Keefer MC, Kallas EG, Corey L, Gorse GJ, Belshe RB, Graham BS, Spearman PW, Schwartz D, Clements ML,

- Mulligan MJ, Goepfert P, Fast PE, Francis D, and the NIAID AIDS Vaccine Evaluation Group. QS-21 promotes an antigen dose-sparing adjuvant effect during HIV-1 envelope subunit immunization in humans. *Vaccine* 2001; (in press).
72. Graham BS, Belshe RB, Clements ML, et al. Vaccination of vaccinia-naive adults with human immunodeficiency virus type 1 gp160 recombinant vaccinia virus in a blinded, controlled, randomized clinical trial. The AIDS Vaccine Clinical Trials Network. *J Infect Dis* 1992; 166:244-52.
73. Cooney EL, McElrath MJ, Corey L, et al. Enhanced immunity to human immunodeficiency virus (HIV) envelope elicited by a combined vaccine regimen consisting of priming with a vaccinia recombinant expressing HIV envelope and boosting with gp160 protein. *Proc Natl Acad Sci USA* 1993; 90:1882-6.
74. Graham BS, Gorse GJ, Schwartz DH, et al. Determinants of antibody response after recombinant gp160 boosting in vaccinia-naive volunteers primed with gp160-recombinant vaccinia virus. *J Infect Dis* 1994; 170:782-6.
75. Pincus SH, Messer KG, Cole R, et al. Vaccine-specific antibody responses induced by HIV-1 envelope subunit vaccines. *J Immunol* 1997; 158:3511-20.
76. Clements-Mann ML, Matthews TJ, Weinhold K, et al. HIV-1 immune responses induced by canarypox (ALVAC)-gp160 MN, SF-2 rgp120, or both vaccines in seronegative adults. *J Infect Dis* 1998; 177:1230-46.
77. Evans TG, Keefer MC, Weinhold KJ, Wolff M, Montefiori DM, Gorse GJ, Graham BS, McElrath MJ, Clements-Mann ML, Mulligan M, Fast PL, Walker MC, Excler JL, Duliege AM, Tartaglia J, and the NIAID AIDS Vaccine Evaluation Group. A canarypox

- vaccine expressing multiple HIV-1 genes given alone or with rgp120 elicits broad and durable CD8⁺ CTL responses in seronegative volunteers. *J Infect Dis* 1999; 180:290-8.
78. Corey L, McElrath MJ, Weinhold K, Matthews T, Stablein D, et al. Cytotoxic T cell and neutralizing antibody responses to HIV-1 envelope with a combination vaccine regimen. *J Infect Dis* 1998; 177:301-9.
 79. Hammond SA, Bollinger RC, Stanhope PE, Quinn TC, Schwartz D, et al. Comparative clonal analysis of human immunodeficiency virus type 1 (HIV-1)-specific CD4⁺ and CD8⁺ cytolytic T lymphocytes isolated from seronegative humans immunized with candidate HIV-1 vaccines. *J Exp Med* 1992; 176:1531-42.
 80. El-Daher N, Keefer MC, Reichman RC, Dolin R, Roberts NJ. Persisting human immunodeficiency virus type 1 gp160-specific human T lymphocyte responses including CD8⁺ cytotoxic activity after receipt of envelope vaccines. *J Infect Dis* 1993; 168:306-13.
 81. Clements-Mann ML, Matthews TJ, Weinhold K, Graham BS, Gorse GJ, et al. HIV-1 immune responses induced by canarypox (ALVAC)-gp160 MN, SF-2 rgp120, or both vaccines in seronegative adults. *J Infect Dis* 1998; 177:1230-46.
 82. Evans TG, Keefer MC, Weinhold KJ, Wolff M, Montefiori DM, et al. A canarypox vaccine expressing multiple HIV-1 genes given alone or with rgp120 elicits broad and durable CD8⁺ CTL responses in seronegative volunteers. *J Infect Dis* 1999; 180:290-8.
 83. Belshe RB, Gorse GJ, Mulligan MJ, Evans TG, Keefer MC, et al. Induction of immune responses to HIV-1 by canarypox virus (ALVAC) HIV-1 and gp120 SF-2 recombinant vaccines in uninfected volunteers. *AIDS* 1998; 12:2407-15.

84. Salmon-Ceron D, Excler JL, Finkielsztejn L, Autran B, Gluckman JC, et al. Safety and immunogenicity of a live recombinant canarypox virus expressing HIV type 1 gp120 MN tm/gag/protease LAI (ALVAC-HIV, vCP205) followed by a p24E-V3 MN synthetic peptide (CLTB-36) administered in healthy volunteers at low risk for HIV infection. *AIDS Res Hum Retrovir* 1999; 15:633-45.
85. Belshe RB, Stevens C, Gorse GJ, Buchbinder S, Weinhold K, et al. Safety and immunogenicity of a canarypox-vectored human immunodeficiency virus type 1 vaccine with or without gp120: a phase 2 study in higher- and lower-risk volunteers. *J Infect Dis* 2001; 183:1343-52.
86. Ferrari G, Humphrey W, McElrath MJ, Excler JL, Duliege AM, et al. Clade B-based HIV-1 vaccines elicit cross-clade cytotoxic T lymphocyte reactivities in uninfected volunteers. *Proc Natl Acad Sci USA* 1997; 94:1396-401.
87. Castillo RC, Arango-Jaramillo S, John R, Weinhold K, Kanki P, et al. Resistance to human immunodeficiency virus type 1 in vitro as a surrogate of vaccine-induced protective immunity. *J Infect Dis* 2000; 181:897-903.
88. Martin SJ, Vyakarnam A, Cheingsongpopov R, et al. Immunization of human HIV-seronegative volunteers with recombinant p17/p24:Ty virus-like particles elicits HIV-1 p24-specific cellular and humoral immune responses. *AIDS* 1993; 7:1315-23.
89. Gahery-Segard H, Pialoux G, Charmeteau B, Sermet S, Poncelet H, Raux M, Tartar A, Levy JP, Gras-Masse H, Guillet JG. Multiepitopic B- and T-cell responses induced in humans by a human immunodeficiency virus type 1 lipopeptide vaccine. *J Virol* 2000; 74:1694-703.

90. Murali-Krishna K, Altman JD, Suresh M, Sourdive DJ, Zajac AJ, et al. 1998. Counting antigen-specific CD8 T cells: a reevaluation of bystander activation during viral infection. *Immunity* 8:177-87.
91. Kovacs JA, Vasudevachari MB, Easter M, et al. Induction of humoral and cell-mediated anti-human immunodeficiency virus (HIV) responses in HIV seronegative volunteers by immunization with recombinant gp160. *J Clin Invest* 1993; 92:919-28.
92. Keefer MC, Graham BS, Belshe RB, et al. Studies of high doses of a human immunodeficiency virus type 1 recombinant glycoprotein 160 candidate vaccine in HIV type 1-seronegative humans. *AIDS Res Hum Retrovir* 1994; 10:1713-23.
93. Wintch J, Chagnat CL, Braun DG, et al. Safety and immunogenicity of a genetically engineered human immunodeficiency virus vaccine. *J Infect Dis* 1991; 163:219-25.
94. Schwartz DH, Gorse G, Clements ML, et al. Induction of HIV-1-neutralising and syncytium-inhibiting antibodies in uninfected recipients of HIV-1IIIB rgp120 subunit vaccine. *Lancet* 1993; 342:69-73.
95. Kahn JO, Sinangil F, Baenziger J, et al. Clinical and immunologic responses to human immunodeficiency virus (HIV) type 1(SF2) gp120 subunit vaccine combined with MF59 adjuvant with or without muramyl tripeptide dipalmitoyl phosphatidylethanolamine in non-HIV-infected human volunteers. *J Infect Dis* 1994; 170:1288-91.
96. Gorse GJ, McElrath MJ, Matthews TJ, et al. Modulation of immunologic responses to HIV-1 recombinant gp160 vaccine by dose and schedule of administration. *Vaccine* 1998; 16:493-506.

97. Gorse GJ, Corey L, Patel GB, Mandava M, Hsieh RH, Matthews TJ, Walker MC, McElrath MJ, Berman PW, Eibl MM, Belshe RB, and the NIAID AIDS Vaccine Evaluation Group. HIV-1MN recombinant glycoprotein 160 vaccine-induced cellular and humoral immunity boosted by HIV-1MN recombinant glycoprotein 120 vaccine. *AIDS Res Hum Retrovir* 1999; 15:115-32.
98. Salmon-Ceron D, Excler JL, Sicard D, et al. Safety and immunogenicity of a recombinant HIV type 1 glycoprotein 160 boosted by a V3 synthetic peptide in HIV-negative volunteers. *AIDS Res Hum Retrovir* 1995; 11:1479-86.
99. Nitayaphan S, Khamboonruang C, Sirisophana N, Morgan P, Chiu J, Duliege AM, Chenchitra C, Supamongse T, Rungreugthanakit K, deSouza M, Mascola JR, Boggio K, Ratto-Kim S, Markowitz LE, Birx D, Suriyanon V, McNeil JG, Brown AE, Michael RA. A phase I/II trial of HIV-1 SF-2 gp120/MF59 vaccine in seronegative Thais. AFRIMS-RIHES Vaccine Evaluation Group. Armed Forces Research Institute of Medical Sciences and the Research Institute for Health Sciences. *Vaccine* 2000;18:1448-55.
100. McCormack S, Tilzey A, Carmichael A, Gotch F, Kepple J, Newberry A, Jones G, Lister S, Beddows S, Cheingsong R, Rees A, Babiker A, Banatvala J, Bruck C, Darbyshire J, Tyrrell D, Van Hoecke C, Weber J. A phase I trial in HIV negative healthy volunteers evaluating the effect of potent adjuvants on immunogenicity of a recombinant gp120W61D derived from dual tropic R5X4 HIV-1ACH320. *Vaccine* 2000; 18:1166-77.
101. Sarin PS, Mora CA, Naylor PH, et al. HIV-1 p17 synthetic peptide vaccine HGP-30: induction of immune response in human subjects and preliminary evidence of protection against HIV challenge in SCID mice. *Cell Mol Biol* 1995; 41:401-7.

102. Kahn JO, Stites DP, Scillian J, et al. A phase I study of HGP-30, a 30 amino acid subunit of the human immunodeficiency virus (HIV) p17 synthetic peptide analogue sub-unit vaccine in seronegative subjects. *AIDS Res Hum Retrovir* 1992; 8:1321-5.
103. Naylor PH, Sztein MB, Wada S, et al. Preclinical and clinical studies on immunogenicity and safety of the HIV-1 p17-based synthetic peptide AIDS vaccine--HGP-30-KLH. *Internatl J Immunopharmacol* 1991; 13 (Suppl) 1:117-27.
104. Gorse GJ, Keefer MC, Belshe RB, et al. A dose-ranging study of a prototype synthetic HIV-1 V3 branched peptide vaccine. *J Infect Dis* 1996; 173:330-9.
105. Li D, Forrest BD, Li Z, et al. International clinical trials of HIV vaccines: II. Phase I trial of an HIV-1 synthetic peptide vaccine evaluating an accelerated immunization schedule in Yunnan, China. *Asian Pacific J All Immunol* 1997; 15:105-13.
106. Kelleher AD, Emery S, Cunningham P, et al. Safety and immunogenicity of UBI HIV-1(MN) octameric V3 peptide vaccine administered by subcutaneous injection. *AIDS Res Hum Retrovir* 1997; 13:29-32.
107. Phanuphak P, Teeratakulpixarn S, Sarangbin S, et al. International clinical trials of HIV vaccines: I. Phase I trial of an HIV-1 synthetic peptide vaccine in Bangkok, Thailand. *Asian Pacific J All Immunol* 1997; 15:41-8.
108. Rubinstein A, Goldstein H, Pettoello-Mantovani M, et al. Safety and immunogenicity of a V3 loop synthetic peptide conjugated to purified protein derivative in HIV-seronegative volunteers. *AIDS* 1995; 9:243-51.
109. Lambert JS, Keefer M, Mulligan MJ, Schwartz D, Mestecky J, Weinhold K, Smith C, Hsieh R, Moldoveanu Z, Fast P, Forrest B, Koff W. A Phase I safety and

- immunogenicity trial of UBI® microparticulate monovalent HIV-1 MN oral peptide immunogen with parenteral boost in HIV-1 seronegative human subjects. *Vaccine* 2001; 19:3033-3042.
110. Boyer JD, Cohen AD, Vogt S, Schumann K, Nath B, Ahn L, Lacy K, Bagarazzi ML, Higgins TJ, Baine Y, Ciccarelli RB, Ginsberg RS, MacGregor RR, Weiner DB. Vaccination of seronegative volunteers with a human immunodeficiency type 1 env/rev DNA vaccine induces antigen-specific proliferation and lymphocyte production of beta-chemokines. *J Infect Dis* 2000; 181:476-83.
 111. Hanke T, McMichael AJ. Design and construction of an experimental HIV-1 vaccine for a year-2000 clinical trial in Kenya. *Nat Med.* 2000; 6:951-5.
 112. Zagury D, Bernard J, Cheynier R, et al. A group specific anamnestic immune reaction against HIV-1 induced by a candidate vaccine against AIDS. *Nature* 1988; 332:728-31.
 113. Cooney EL, Collier AC, Greenberg PD, et al. Safety of and immunological response to a recombinant vaccinia virus vaccine expressing HIV envelope glycoprotein. *Lancet* 1991; 337:567-72.
 114. Pialoux G, Excler JL, Riviere Y, et al. A prime-boost approach to HIV preventive vaccine using a recombinant canarypox virus expressing glycoprotein 160 (MN) followed by a recombinant glycoprotein 160 (MN/LAI). *AIDS Res Hum Retrovir* 1995; 11:373-81.
 115. Fleury B, Janvier G, Pialoux G, et al. Memory cytotoxic T lymphocyte responses in human immunodeficiency virus type I (HIV-1)-negative volunteers immunized with a recombinant canarypox expressing gp160 of HIV-1 and boosted with a recombinant gp160. *J Infect Dis* 1996; 174:734-8.

116. Belshe RB, Gorse GJ, Mulligan MJ, Evans TG, Keefer MC, Excler JL, Duliege AM, Tartaglia J, Cox WI, McNamara J, Hwang KL, Bradney A, Montefiori D, Weinhold KJ. Induction of immune responses to HIV-1 by canarypox virus (ALVAC) HIV-1 and gp120 SF-2 recombinant vaccines in uninfected volunteers. NIAID AIDS Vaccine Evaluation Group. *AIDS* 1998; 12:2407-15.
117. Salmon-Ceron D, Excler JL, Finkielsztejn L, Autran B, Gluckman JC, Sicard D, Matthews TJ, Meignier B, Valentin C, El Habib R, Blondeau C, Raux M, Moog C, Tartaglia J, Chong P, Klein M, Milcamps B, Heshmati F, Plotkin S. Safety and immunogenicity of a live recombinant canarypox virus expressing HIV type 1 gp120 MN MN tm/gag/protease LAI (ALVAC-HIV, vCP205) followed by a p24E-V3 MN synthetic peptide (CLTB-36) administered in healthy volunteers at low risk for HIV infection. AGIS Group and L'Agence Nationale de Recherches sur Le Sida. *AIDS Res Hum Retroviruses* 1999; 15:633-45.
118. McElrath MJ, Corey L, Montefiori D, Wolff M, Schwartz D, Keefer M, Belshe R, Graham BS, Matthews T, Wright P, Gorse G, Dolin R, Berman P, Francis D, Duliege AM, Bolognesi D, Stablein D, Ketter N, Fast P. A Phase II study of two HIV type 1 envelope vaccines, comparing their immunogenicity in populations at risk for acquiring HIV type 1 infection. AIDS Vaccine Evaluation Group. *AIDS Res Hum Retroviruses* 2000; 16:907-19.
119. Belshe RB, Stevens C, Gorse GJ, Buchbinder S, Weinhold K, Sheppard H, Stablein D, Self S, McNamara J, Frey S, Flores J, Excler JL, Klein M, Habib RE, Duliege AM, Harro C, Corey L, Keefer M, Mulligan M, Wright P, Celum C, Judson F, Mayer K, McKirnan

D, Marmor M, Woody G. Safety and immunogenicity of a canarypox-vectored human immunodeficiency virus Type 1 vaccine with or without gp120: a phase 2 study in higher- and lower-risk volunteers. *J Infect Dis* 2001; 183:1343-52.